

Topical management of tinea pedis

Betsy Thomas BScPharm Jamison Falk PharmD G. Michael Allan MD CCFP

Clinical question

How effective are topical treatments for tinea pedis?

Bottom line

Tinea pedis is successfully treated with topical antifungals in 70% to 75% of patients compared with 20% to 30% using placebo. Tea tree oil is likely ineffective. Topical terbinafine might result in an absolute improvement of 2% to 8% more patients cured over other topicals. Most patients were treated for 1 week with terbinafine and 4 to 6 weeks with azoles (like clotrimazole).

Evidence

Results are statistically significant unless indicated.

- Studies comparing with placebo found the following.
 - Systematic review (67 RCTs) of mycologically diagnosed tinea pedis (and onychomycosis, not included here), reporting laboratory-confirmed treatment failure at 6 weeks.¹
 - Allylamines (eg, terbinafine, naftifine), 9 RCTs (N=876), 1 to 4 (most 4) weeks' treatment: 25% versus 80% placebo; number needed to treat (NNT) of 2.
 - Azoles (eg, clotrimazole, miconazole), 6 RCTs (N=448), 4 to 6 weeks' treatment: 28% versus 70% placebo; NNT=3.
 - Tea tree oil, 1 RCT (N=71), 4 weeks' treatment: no difference from placebo.
 - Systematic review of topical terbinafine versus placebo, 9 RCTs (N=986), 1 to 4 (1 most common) weeks' treatment²: clinical cure, 72% terbinafine versus 28% placebo; NNT=3.
 - Other systematic reviews found similar results.³⁻⁵
- Direct comparisons found the following.
 - Allylamines versus azoles.
 - Systematic review, 8 RCTs (N=1034), 1 to 6 (most 1 to 2) weeks' treatment³: mycological cure, 78% allylamines versus 76% azoles; NNT=40.
 - Topical terbinafine versus other antifungals.
 - Systematic review, 10 RCTs (N=1341), 1 to 4 (most 1) weeks' treatment²: clinical cure, 83% terbinafine versus 75% other antifungals (statistical significance reported inconsistently; if real, NNT=13).
- Adverse events: burning, stinging, and itching sensations were most common (but not quantified).¹
- Limitations: some RCTs¹ and 1 systematic review² were industry funded; clinical cure is less commonly reported than mycological cure.

Context

- Topical antifungals are suggested as first-line agents, reserving oral agents for severe disease (eg, moccasin-type infection), failed topical treatment, and immunocompromised patients.⁶
- Approximate cost (for 30 g)⁷⁻⁹:
 - 1% clotrimazole cream, twice daily, \$13,
 - 2% miconazole cream, twice daily, \$15, and
 - 1% terbinafine cream, twice daily, \$30.

Implementation

It is uncertain whether foot hygiene or changing footwear is beneficial; however, placebo arms from RCTs suggest it might help.⁴ The Centers for Disease Control and Prevention advise that patients with tinea pedis keep feet dry, clean, and cool; wear sandals, if possible (especially in locker rooms); air out shoes; and wear cotton socks.¹⁰ Patients can discuss over-the-counter options with their pharmacists, such as clotrimazole or miconazole, although these tend to require a longer treatment duration. Nystatin should not be used owing to dermatophyte resistance.⁶ Terbinafine cream, twice a day for 7 days, is a reasonable prescription option with a short treatment duration and well supported efficacy. 🌿

Ms Thomas is Clinical Evidence Expert at the College of Family Physicians of Canada. **Dr Falk** is Associate Professor in the College of Pharmacy at the University of Manitoba in Winnipeg. **Dr Allan** is Director of Programs and Practice Support at the College of Family Physicians of Canada.

Competing interests

None declared

The opinions expressed in Tools for Practice articles are those of the authors and do not necessarily mirror the perspective and policy of the Alberta College of Family Physicians.

References

- Crawford F, Hollis S. Topical treatments for fungal infections of the skin and nails of the foot. *Cochrane Database Syst Rev* 2007;(3):CD001434.
- Korting HC, Kiencke P, Nelles S, Rychlick R. Comparable efficacy and safety of various topical formulations of terbinafine in tinea pedis irrespective of the treatment regimen: results of a meta-analysis. *Am J Clin Dermatol* 2007;8(6):357-64.
- Rotta I, Otuki MF, Sanches AC, Correr CJ. Efficacy of topical antifungal drugs in different dermatomycoses: a systematic review with meta-analysis. *Rev Assoc Med Bras* (1992) 2012;58(3):308-18.
- Crawford F. Athlete's foot. *BMJ Clin Evid* 2009;2009:1712.
- Rotta I, Sanchez A, Gonçalves PR, Otuki MF, Correr CJ. Efficacy and safety of topical antifungals in the treatment of dermatomycosis: a systematic review. *Br J Dermatol* 2012;166(5):927-33.
- Ely JW, Rosenfeld S, Seabury Stone M. Diagnosis and management of tinea infections. *Am Fam Physician* 2014;90(10):702-10.
- Interactive drug benefit list. Alberta Health; 2020. Available from: <https://idbl.ab.bluecross.ca/idbl/load.do>. Accessed 2020 Mar 21.
- Micatin cream 2% - miconazole nitrate cream USP, 30 g. Mississauga, ON: Walmart Corporation; 2020. Available from: <https://www.walmart.ca/en/ip/micatin-cream-2-miconazole-nitrate-cream-usp-30-g/6000189068419>. Accessed 2020 Jul 13.
- Canesten 1% topical antifungal cream. Mississauga, ON: Walmart Corporation; 2020. Available from: <https://www.walmart.ca/en/ip/bayer-healthcare-consumer-care-canesten-1-topical-antifungal-cream/6000017348217>. Accessed 2020 Jul 13.
- Hygiene-related disease. Athlete's foot (tinea pedis). Bethesda, MD: Centers for Disease Control and Prevention; 2017. Available from: https://www.cdc.gov/healthywater/hygiene/disease/athletes_foot.html. Accessed 2020 Oct 26.

Can Fam Physician 2021;67:30. DOI: 10.46747/cfp.670130

Tools for Practice articles in *Canadian Family Physician* are adapted from articles published on the Alberta College of Family Physicians (ACFP) website, summarizing medical evidence with a focus on topical issues and practice-modifying information. The ACFP summaries and the series in *Canadian Family Physician* are coordinated by Dr G. Michael Allan, and the summaries are co-authored by at least 1 practising family physician and are peer reviewed. Feedback is welcome and can be sent to toolsforpractice@cfpc.ca. Archived articles are available on the ACFP website: www.acfp.ca.